

Ionized gas, molecules, and dust in Sh2-132

J. Vasquez^{1,2*}, C.E. Cappa^{1,2}, S. Pineault^{1,3} and N.U. Duronea^{1,2}

¹*Instituto Argentino de Radioastronomía, CCT-La Plata, CONICET, C.C.5., 1894, Villa Elisa, Argentina*

²*Facultad de Ciencias Astronómicas y Geofísicas, Universidad Nacional de La Plata, La Plata, Argentina*

³*Département de physique, de génie physique et d'optique and Centre de recherche en astrophysique du Québec (CRAQ), Université Laval, Québec, Canada G1VOA6*

Accepted 1988 December 15. Received 1988 December 14; in original form 1988 October 11

ABSTRACT

We analyze the various interstellar components of the HII region Sh2-132. The main stellar source is the double binary system that includes the Wolf-Rayet star WR153ab. We use radio continuum images at 408 and 1420 MHz, and H I 21cm line data taken from the Canadian Galactic Plane Survey, molecular observations of the ¹²CO(1-0) line at 115 GHz from the Five College Radio Astronomy Observatory, and available mid and far IR observations obtained with the MSX and IRAS satellites, respectively.

Sh2-132 is composed of two shells showing radio continuum counterparts at both frequencies. The emission is thermal in nature. The estimated rms electron density and ionized mass of the nebula are $n_e \simeq 20 \text{ cm}^{-3}$ and $M_{\text{HII}} \simeq 1500 M_{\odot}$. The distribution of the CO emission shows molecular gas bordering the ionized nebula and interacting with it. The velocities of the molecular gas is in the range -38 to -53 km s^{-1} , similar to the velocity of the ionized gas.

The emission at $8.3 \mu\text{m}$ reveals a ring like feature of about $15'$ that encircles the bright optical regions. This emission is due to the PAHs and marks the location of photodissociation regions.

The gas distribution in the environs of Sh2-132 can be explained in a scenario where the massive stars in the region photodissociated, ionized, and swept-up the dense molecular material from the parental cloud through their strong stellar winds and intense UV photon flux.

Key words: ISM: bubbles – ISM: molecules – ISM: HII regions – ISM: individual: Sh2-132 – stars: Wolf-Rayet – stars: individual: WR 153ab

1 INTRODUCTION

The interaction between massive stars and the interstellar medium leads to the formation of HII regions, interstellar bubbles, and new stars. More specifically, the interaction between the molecular medium and the UV stellar radiation is at the origin of photodissociation regions (PDRs). Molecules are dissociated by UV photons in the range 6 eV to 13.6 eV. The gas component ranges from ionized atoms at the PDR surface to molecules deeper into the molecular region as the FUV flux is absorbed. The FUV flux illuminating these regions range between $1 \leq G_0 \leq 10^6$, where G_0 is the intensity of the radiation field in the solar neighborhood (Habing 1968), while the H I density lies between $10 \text{ cm}^{-3} \leq n_H \leq 10^5 \text{ cm}^{-3}$ (Tielens & Hollenbach 1985, Sternberg & Dalgarno 1989, Sternberg & Dalgarno 1995, Bakes & Tielens

1998). PDRs are formed at the edge of HII regions, planetary nebulae, reflexion nebulae, and stellar forming regions. Forbidden spectral lines in the far infrared as well as molecular lines are observed towards PDRs (e.g. Hogerheijde et al. 1995, Kaufman et al. 1999).

An important component of PDRs is interstellar dust. Its presence plays a major role absorbing stellar radiation and reemitting it in the mid and far infrared. Polycyclic aromatic hydrocarbons (PAHs), which emit at 3.3, 6.2, 7.7, 8.6, and $11.3 \mu\text{m}$, are tracers of PDRs (e.g. Watson et al. 2008).

In this paper, we study the interstellar environment of the HII region Sh2-132 (Sharpless 1959), located in the Perseus spiral arm, paying particular attention to the presence of PDRs at the interfase between the ionized and molecular gas.

Sh2-132 was detected in H α , [O III], and [S II] lines by Heckathorn et al. (1982). The nebula is centered approxi-

* E-mail: jvasquez@fcaglp.unlp.edu.ar; pete@iar-conicet.gov.ar

mately at $(l, b) \simeq (102^\circ 50', -0^\circ 42')$ and presents two bright regions, which are surrounded by diffuse emission regions. They can be identified in Fig. 1, which displays the DSS-R image of the nebula.

Churchwell & Walmsley (1973) performed radio continuum observations at 2695 MHz with an angular resolution of $18''.2$, and, using measurements at several frequencies, derived a thermal spectral index. Harten et al. (1978) investigated the characteristics and morphology of the nebula based on radio continuum observations at 610 MHz using the WSRT with a synthesized beam of $60''$. Most of the emission at this frequency originates from two optically bright regions named Shell A and Shell B, located at $(l, b) \simeq (102^\circ 57', -0^\circ 42')$ and $(l, b) \simeq (102^\circ 47', -0^\circ 42')$. Both shells coincide with the bright optical regions within the nebula and are indicated in Fig. 1.

According to Harten et al. (1978), the optically bright section of Sh2-132 (at $l > 102^\circ 43'$) is an ordinary HII region excited by the massive O-type star BD+55°2722 and the WR star WR 153ab (= HD 211853), which are located close to the center of the nebula and linked to Shell B. Chu et al. (1983) reached to similar conclusions, based on the presence of the stars near the center of the nebula and on the absence of velocity changes in the nebular spectra. The main parameters of these stars, as well as of other OB stars listed by Harten et al. (1978) that appear projected onto the HII region, are summarized in Table 1, which lists their galactic coordinates, spectral classification, visual magnitude V , colour indices $(B-V)$ and $(B-V)_0$, visual absorption A_V , absolute magnitude M_V , and derived spectrophotometric distance d . Uncertainties in the derived spectrophotometric distances of some of the stars were estimated using M_V values from Vacca et al. (1996) and Landolt-Börnstein (1982). The large distance derive for LS III+55°37 casts doubts on their relation to the nebula. Note that, based on photometric data, Panov & Seggewiss (1990) concluded that HD 211853 is a system with two pairs of stars, both hosting a WR component and an O-type star.

Chu & Treffers (1981), based on Fabry Perot observations of the $H\alpha$ line, found that the ionized gas has LSR velocities in the range -53 to -45 km s^{-1} , in agreement with previous results from Georgelin & Georgelin (1976, -50.9 km s^{-1}), Reynolds (1988, -48 ± 1 km s^{-1}), and Fich et al. (1990, -47.1 km s^{-1}). Quireza et al. (2006) detected ionized gas at -50.46 and -49.48 km s^{-1} from He and H radio recombination lines, respectively. Harten et al. (1978) estimated an emission measure $EM = 3000$ pc cm^{-6} for the entire nebula.

New $H\alpha$ and [O III] images taken by Miller & Chu (1993) revealed an arc of $4'$ in size to the south of the WR star (at $[l, b] = [102^\circ 46', -0^\circ 43']$). They concluded that the filament probably indicates the position of an ionization front and considered it to be a ring nebula related to the star. Based on optical data, Esteban & Rosado (1995) found velocities in the range -41 to -59 km s^{-1} for the arc and derived an electron density of 290 ± 100 cm^{-3} from line ratios. The [OIII]/ $H\alpha$ ratio is consistent with values derived for ring nebulae, suggesting that WR 153ab is the main source of ionization (Esteban & Rosado 1995). These authors classify the arc as R_s -type based on the morphology of this feature and on the absence of a clear evidence of expanding motions in the line profiles. This last point suggests that the massive




Figure 1. DSS-R image of the brightest section of Sh2-132. The grey scale is arbitrary. The large cross marks the position of the WR star, while smaller crosses indicate the location of the OB stars.

progenitor of the WR star contributed in the shaping of the nebula (see Chu et al. 1981).

Distance estimates for the WR star based on different methods range from 2.75 to 5.0 kpc. Massey (1981) derived a spectrophotometric distance of 5.0 kpc, while van der Hucht (2001) estimated 2.75 kpc based on its association to Cep OB1, with a distance uncertainty of 40%. Foster & Routledge (2003), using a method based on HI column densities, estimated 3.2 ± 0.5 kpc. A distance of 3.68 kpc was adopted by Churchwell & Wamsley (1973), based on the WN6+BOI spectral type of the WR star, known at that time. Georgelin & Georgelin (1976) derived a distance of 3.6 kpc, while Fich, Blitz & Stark (1989) obtained 4.2 ± 1.5 kpc.

Adopting a mean LSR velocity of -48 km s^{-1} for the ionized gas, and taking into account the presence of non-circular motions in this section of the Galaxy (see Fig. 2a by Brand & Blitz 1993), a kinematical distance $d_k = 3.5 \pm 1.0$ kpc can be predicted, in close agreement with distances derived optically for the WR star. We adopt this distance in the present study.

With the aim of investigating in some detail the nature of Sh2-132, we analyzed the distribution of the different gas components in its environs. To accomplish this goal we performed a multi-frequency study using radio continuum data at several frequencies, HI 21cm line data, $^{12}\text{CO}(1-0)$ line data, and IR observations at different wavelengths. These data revealed that photodissociation regions are present at the interface between the ionized and molecular gas.

2 DATABASE

Neutral hydrogen 21cm line data from the Canadian Galactic Plane Survey (CGPS, Taylor et al. 2003) was used to analyze the HI gas distribution, while $^{12}\text{CO}(1-0)$ data at 115 GHz from the Five College Radio Astronomy Observatory, allowed us to study the molecular distribution. Intensities are shown as main beam brightness temperatures T_{mb} . The

Table 1. O-B type stars towards the bright section of Sh2-132

Star	(<i>l</i> , <i>b</i>)	ST	<i>V</i> mag	(<i>B</i> − <i>V</i>) mag	(<i>B</i> − <i>V</i>) ₀ mag	<i>A_v</i> mag	<i>M_V</i> mag	<i>d</i> kpc
HD 211853 (WR 153ab)	102°46′8, −0°39′	WN6/WC+O6I ⁽¹⁾	9.08 ⁽²⁾	0.27 ⁽²⁾	(2)	2.28 ⁽²⁾	−6.4 ⁽²⁾	2.75 ⁽²⁾ , 5.0 ⁽⁴⁾
LS III+55°37	102°46′8, −0°40′2	O7V ⁽³⁾	11.01	0.32 ⁽³⁾	−0.29 ⁽⁶⁾	1.86	−5.2 ⁽⁵⁾	7.5
BD+55°2722	102°48′6, −0°40′2	O8.5V ⁽³⁾	9.91 ⁽³⁾	0.45 ⁽³⁾	−0.28 ⁽⁶⁾	2.26	−4.7 ⁽⁵⁾	3.0
LS III+55°40	102°56′4, −0°42′6	O9V ⁽³⁾	12.08 ⁽³⁾	0.64 ⁽³⁾	−0.28 ⁽⁶⁾	2.85	−4.5 ⁽⁵⁾	5.6
LS III+55°42	102°56′4, −0°43′8	B ⁽⁷⁾	12.38	0.52	−0.26 ⁽⁶⁾	2.42	−2.45, −5.1 ^(**5)	5-10-13
LS III+55°39	102°57′6, −0°38′4	B ⁽⁷⁾	11.74	0.45	−0.26 ⁽⁶⁾	2.20	−2.45, −5.1 ^(**5)	2.5-8.5
LS III+55°45	102°59′4, −0°45′6	O8 ⁽³⁾	10.45 ⁽³⁾	0.70 ⁽³⁾	−0.27 ⁽⁵⁾	3.01	−4.9, −5.8 ^(**5)	2.9-4.9
LS III+55°49	103°02′4, −0°48′6	B ⁽⁷⁾	11.50	0.57	−0.26 ⁽⁶⁾	2.5 7	−2.45, −5.1 ^(**5)	1.9-6.4

References: (1) Smith et al. (1998), (2) van der Hucht (2001), (3) GOS Catalog V2.2, Sota et al. (2008), (4) Massey (1981), (5) Landolt-Rörmstein (1996), (6) Wegner 1994, (7) SIMBAD database, (*) Spectral classification B0III-B2V was adopted, (**) spectral class V and III was adopted.

Table 2. Observational parameters of the H I and ¹²CO data

	H I	¹² CO(1-0)
Original FWHM	1′18×0′98	~1′
Final FWHM	1′4×1′4	
Vel. range (LSR) (km s ^{−1})	−165, +57	−165, +57
Vel. resolution (km s ^{−1})	1.32	0.824
RMS noise <i>T_b</i> (K)	1.7	0.15

H I data were smoothed to 1′4 × 1′4 to facilitate the identification of the H I features. The main observational parameters of these data bases are detailed in Table 2.

Radio continuum observations at 408 and 1420 MHz were also extracted from the CGPS. The image at 408 MHz was obtained with a synthesized beam and an rms noise of 3′4×2′8 and ~1.1 K, respectively; the corresponding values for 1420 MHz are 1′0×0′81, and 0.063 K.

Mid-infrared (MIR) images were extracted from the MSX Galactic Plane Survey (Price et al. 2001) at 8.3 μm (A-band), 12.1 μm (C-band), 14.7 μm (D-band), and 21.3 μm (E-band), with an angular resolution of 18′′.4. MIR and far-infrared (FIR) images at 12, 25, 60, and 100 μm from the IRAS satellite (HIRES data) are also included. These images have angular resolutions in the range 0′5 to 2′0.

3 RESULTS

3.1 Radio continuum emission

Figures 2 and 3 show the CGPS radio continuum emission at 1420 MHz. The grayscale in Fig. 2 has been chosen to emphasize the large-scale region of faint emissivity to the north of WR 152 (Cappa et al. 2010, hence the “burnt-out” appearance of the region around WR 153ab). The emission distribution at 1420 MHz resembles an upside-down “U”. This semicircular structure consists of two areas of strong emission coincident with Shells A and B detected by Harten et al. (1978), centered at (*l*, *b*) ≃ (102°34′42′′.4, −0°46′9′′.8) and at (*l*, *b*) ≃ (102°26′35′′.8, −0°43′10′′.2), respectively. WR 153ab, indicated by the larger cross in Fig. 1, appears projected onto the central region of Shell B. Thus, the bright radio




Figure 2. Full resolution CGPS image at 1420 MHz emphasizing faint emission to the north of WR 152 (indicated by a cross symbol in the lower right part of the figure). The arrow points to a continuum feature corresponding to the large HI structure.

emission at *l* ≥ 102°5 is in excellent agreement with the optical emission (see Fig. 1) and with the image at 610 MHz by Harten et al. (1978). Both Shells A and B can be identified at 1420 MHz. Note in particular the small arc of emission at (*l*, *b*) = (102°77, −0°73) (best identified in Fig. 3), which is the radio counterpart of the 4′-arc detected at optical wavelengths by Miller & Chu (1993).

The faint structure centered at (*l*, *b*) = (102°2, −0°83) is related to WR 152 and has been analyzed in a previous paper (Cappa et al. 2010). Of particular interest is the curved filament indicated by an arrow in Fig. 2. This filament can be followed from *b* ≈ −0°75 to nearly *b* ≈ −0°05. The figure also shows a faint barely visible northeastern section. This structure is likely the radio continuum counterpart of the ~60′ diameter H I structure discussed in Cappa et al. (2010).

Three bright point sources, best seen on Figure 3, are




Figure 3. Full resolution 1420 MHz CGPS image showing the eastern and western regions of Sh2-132 used for TT-plot analyses. The radio sources in direction to Sh2-132 are indicated by 10, 13 and 14. The white contour line marks the 9 K contour of the convolved 1420 MHz image used for the analysis. See text for details.

projected onto the nebula near Shell A. Their coordinates, flux densities and identifications are included in Table 3. The spectral index estimates indicate that sources 10 and 14 are non-thermal in nature and probably extragalactic, while source 13 is thermal. The value of the spectral index indicates that the radio source is optically thick at least at 408 MHz.

The CGPS image at 408 MHz does not show as many details as the one at 1420 MHz, as it is of much lower resolution. However it can be used to study the spectral index distribution of the radio continuum emission around WR 153ab. As a first step, we removed the point sources which are present over the region of interest and convolved the 1420 and 408 MHz images to the same spatial resolution of $3''.4$. We then constructed a TT-plot, in which the brightness temperature T_b at one frequency is plotted point-by-point against the brightness temperature at the other frequency. The brightness temperature spectral index β , where $T_b \propto \nu^\beta$, is directly related to the slope of the straight line fitted by regression, $\beta = -\log(\text{slope})/\log(1420/408)$. The usual flux density spectral index α ($S_\nu \propto \nu^\alpha$) is simply $\alpha = \beta + 2$. The TT-plot method is immune from errors in the zero-level of the individual images (which the simpler technique of dividing one map by the other, even after background subtraction, is not).

Three different TT-plots were constructed, one for the entire Sh2-132 region and one for the eastern and western halves, which correspond to the two shells identified by Harten et al. (1978). Figure 3, which is the full resolution image at 1420 MHz, shows these different regions, the dashed lines delimiting the eastern and western parts. The three bright small diameter sources coincide with sources 10, 13 and 14 in Table 2 of Harten et al. (1978) and had been subtracted from the original image before convolution to a circular beam of $3''.4$.

The results of the TT-plot analysis are presented in




Figure 4. TT-plots corresponding to (a) the whole structure, (b) the eastern half and (c) the western half. For all plots, a background of 9 K and 80 K has been subtracted from the convolved images at 1420 and 408 MHz, respectively.

Figure 4. In each case, the analysis was limited to regions brighter than 9 K at 1420 MHz and 80 K at 408 MHz. Although a significant amount of scatter is present on the diagram, especially for the eastern sector, there is no evidence for the presence of significant variations of the spectral index and a further study of spatial variations is not warranted. The best-fit spectral index is $\alpha = -0.03 \pm 0.03$, corresponding to thermal emission.

In order to obtain some radio continuum parameters, we first derive the total number N_u of UV ionizing photons per second required to produce the observed free-free emission. This can be written as (e.g. Chaisson 1976):

$$N_u = 0.76 \times 10^{47} T_4^{-0.45} \nu_{\text{GHz}}^{0.1} S_\nu d_{\text{kpc}}^2, \quad (1)$$

where S_ν is the flux density in Jy, T_4 is the electron temperature in units of 10^4 K and ν_{GHz} the frequency in GHz. We use the CGPS image at 1420 MHz. The derived flux density is significantly dependent on how the source dimension is defined. The surrounding background is somewhat variable, nevertheless it is reasonable to assume that the source emission corresponds to brightness temperatures above 6.5 to 7 K. Using an average background of 7 K, we obtain a flux density at 1420 MHz of 23 ± 5 Jy, the error being determined from a second estimate using a background of 6.5 K. From the above equation for N_u and taking $T_4 = 0.8$ (Quireza et al. 2006), we obtain $N_u = 2.2 \times 10^{49} \text{ s}^{-1}$. Considering two O6I-type stars and two WN6 as components of the multiple system, and the spectral type from BD+55°2722, LS III+55°45, and LS+55°40 (see Table 1, related to the HII region), we obtain the UV ionizing photons per second emitted by the stars, $Q_0 = (16.6 \pm 3.0) \times 10^{49} \text{ s}^{-1}$, for solar metallicity from Smith et al. (2002).

The comparison between the number of UV photons emitted by the stars Q_0 and the number of UV photons used to ionize the gas N_u indicates that the massive stars in the nebula are responsible for the ionization. A large fraction of the UV photons are also used to heat the associated interstellar dust. This result differs from the one by Harten et al. (1978) since their estimate of Q_0 was too low.

We can now use the models of Mezger & Henderson (1967) to infer the properties of the extended ionized gas, namely its mass M_{HII} and the rms electron number density n_e . Letting θ_{Ga} be the observed half-power width of the extended emission (in minutes of arc, after correction for the beam), we obtain:

$$n_e / \text{cm}^{-3} = 6.35 \times 10^2 u_1 a^{-1/2} T_4^{0.175} \nu_{\text{GHz}}^{0.05} S_\nu^{0.5} \theta_{\text{Ga}}^{-1.5} d_{\text{kpc}}^{-0.5} \quad (2)$$

$$M_{\text{HII}} / M_\odot = 0.386 u_2 a^{-1/2} T_4^{0.175} \nu_{\text{GHz}}^{0.05} S_\nu^{0.5} \theta_{\text{Ga}}^{1.5} d_{\text{kpc}}^{2.5} \quad (3)$$

where $a(\nu, T_e)$ is the Gaunt factor (about unity for our purposes), d_{kpc} is the distance in kpc and the parameters u_1 and u_2 , both of order unity, depend on the assumed model for the source. These parameters are given by Mezger & Henderson (1967) for three different idealized models. From Figure 3, it is apparent that the nebulosity can be roughly broken up into two different regions separated by a nearly vertical line running at a longitude of $\sim 102^\circ 87$. These two regions correspond to shells A and B from Harten et al. (1978). The flux densities of the two sub-regions are 11 ± 2 Jy and 12 ± 3 Jy, respectively. For simplicity, we model each half-region as a sphere of constant density and of angular size $12'$. From Mezger & Henderson (1967), we find $u_1 = 0.78$, $u_2 = 1.29$.




Figure 5. *Upper panel* Sh2-132 at $8.3 \mu\text{m}$ (MSX band A). The grayscale goes from 8.0×10^{-7} to $3.5 \times 10^{-6} \text{ W m}^{-2} \text{ ster}^{-1}$. Extended filaments and point-like sources are indicated (see text). *Bottom panel:* Overlay of the DSS-R image and the image at $60 \mu\text{m}$. IR contours are 50 to $190 \text{ MJy ster}^{-1}$ in steps of 20 MJy ster^{-1} , and from 300 to $900 \text{ MJy ster}^{-1}$ in steps of $200 \text{ MJy ster}^{-1}$.

Assuming $T_4 = 0.8$, from which we determine $a^{-1/2} = 1.04$, we finally have an rms $n_e = 20 \text{ cm}^{-3}$ and $M_{\text{HII}} = 1500 M_\odot$ for each region, where, given the error in estimating fluxes, we have simply taken $S_\nu = 10$ Jy for each half. The derived rms electron density is compatible with estimates by Harten et al. (1978) for Shells A and B.

3.2 IRAS and MSX emission

The IRAS ($60 \mu\text{m}$) and MSX band A ($8.3 \mu\text{m}$) emissions are shown in Fig. 5. Although we analyzed the four IRAS bands, we include the $60 \mu\text{m}$ -image only since the extended emission is clearly seen at this wavelength. Emission at $8.3 \mu\text{m}$ (upper panel) in diffuse nebulae has a contribution from both PAH and dust continuum. As pointed out by Povich et al. (2007), in very energetic regions like M17, dust continuum can be very strong. However, for the case of Sh2-132, the assumption that MSX band A is dominated by PAH emission seems to be reasonable. Consequently, emission in this band indicate the presence of photodissociation regions, while the emission in the far IR originates in large dust grains.

Table 3. Small diameter radio sources towards Sh2-132.

#	l b	S_{1420} mJy	S_{408} mJy	α	S_{178} mJy	S_{365} mJy	S_{610} mJy	S_{1420} mJy	S_{4850}	Identification
(1)		(2)	(2)	(2)	(3)	(4)	(5)	(6)	(5)	
10	102°59'50''90 -0°40'23''35	195±10	550	-0.8		819±87	330		<40	7C 2218+558 ⁽⁷⁾
13	103°3'30''24 -0°41'28''14	365±15	280	+0.2			160	222.6±7.5	<40	NVSS J222034+561438
14	103°4'40''04 -0°47'27''04	450±20	1200	-0.8	3100±500	1738±62	810		60	TXS 2219+559

References: (1) Number from the list of Harten et al. (1978); (2) this paper, derived using CGPS data; (3) Gower et al. (1967); (4) Douglas et al. (1996); (5) Harten et al. (1978); (6) Condon et al. (1998); (7) Waldrum et al. (1996).

The image at 8.3 μm (upper panel) reveals the existence of a ring-like feature centered at $(l, b) = (102^\circ 53', -00^\circ 40')$ of about $\sim 15'$ in size. This feature can be also identified at 60 μm (lower panel). The comparison of the optical emission with the emission at 8.3 μm shows that the mid IR emission encircles most of the nebula, following the outer rim of the ionized gas. This fact is more noticeable for Shell B and is particularly striking near $(l, b) = (102^\circ 47', -00^\circ 43')$, where the 4' ring nebula is present. The bottom panel shows that the emission distribution at 60 μm resembles that in the optical.

In order to estimate some dust parameters, we have divided this ring-like structure in three main filaments, named f1, f2, and f3, which are indicated in the upper panel of Fig. 5. Note that f3 borders the region with high extinction seen in the optical image near $(l, b) = (103^\circ 0', -00^\circ 47')$.

A number of interesting point-like sources located onto the filaments can also be identified in the MSX and IRAS images. They are indicated in the figure as ps1, ps2, and ps3. Infrared source ps1 at $(l, b) = (102^\circ 39'45'', -00^\circ 36'24'')$ is the MSX source G102.6627-00.6067, coincident with IRAS 22160+5551; ps2 at $(l, b) = (102^\circ 48'18'', -00^\circ 43'6'')$ corresponds to G102.8051-00.7184 and IRAS 22172+5549; while ps3 at $(l, b) = (103^\circ 03'44'', -00^\circ 41'27'')$ is G103.0624-00.6911 and IRAS 22187+5559.

After background subtraction, the IR flux densities of the filaments in the four IRAS bands and in the MSX band A were calculated. These values are listed in Table 4. The quoted errors reflect the uncertainties in the determination of the background. Emission in bands C, D, and E is very weak in comparison with the strong background emission leading to large uncertainties in flux density estimates. Values for these bands are not included. The three point sources, whose flux densities in the MSX and IRAS bands are included in the table, can be classified as YSO candidates following criteria by Junkes et al. (1992) and Lumsden et al. (2002) for IRAS and MSX point sources, respectively. In the case of the IRAS criteria, Junkes et al. (1992) consider (i) $S_{100} > 20$ Jy, (ii) $1.2 < S_{100}/S_{60} < 6$, (iii) $S_{60}/S_{25} \geq 1$, and $Q_{60} + Q_{100} \geq 4$, where S_i is the flux density in the bands centered at 25, 60 and 100 μm , and Q_i is the quality of the fluxes at these bands. Lumsden et al.'s (2002) criteria are (i) $S_{21}/S_8 > 2$ and (ii) $S_{14}/S_{12} > 1$, where S_i is the flux density centered at 8.3, 12.1, 14.7, and 21.3 μm .

It has been demonstrated (Scoville & Kwan 1976) that the infrared spectrum of radiatively excited dust grains can be fitted by a modified Planck function $B(T_d)\nu^m$, where ν^m is

the grain emissivity and m increases from 1 to 2 depending on the wavelength and composition of dust grains (Schwartz 1982). Adopting $m = 1.5$, according with the molecular environment (Whittet, 1992), and using the measured flux densities at 60 and 100 μm , we derived the dust color temperature T_d summarized in Table 4. IR luminosities for the point sources are also included. Infrared filaments are excited by the intense stellar radiation field of WR 153ab and the other massive stars in the region. The slightly higher dust color temperature derived for f1 can be explained bearing in mind that this filament is closer to the exciting stars than f2 and f3. In Fig. 6, the MIR and FIR spectral energy distribution corresponding to the filaments are shown. Note that the measured fluxes at shorter wavelengths exceed the values of our model, possibly as the result of the emission of nebular lines which may contribute to the infrared flux densities.

3.3 Molecular gas distribution


The analysis of the $^{12}\text{CO}(1-0)$ emission distribution leads to the detection of two structures with velocities in the ranges from -53.4 to 46.0 km s^{-1} , and from -46.0 to -38.6 km s^{-1} . Fig. 7 shows the averaged CO emission (contours) for the two molecular structures superposed onto the emission at 8.3 μm .

The CO feature present between -46.0 and -38.6 km s^{-1} (upper panel) exhibits a ring-like appearance between $l \sim 102^\circ 35'$ and $102^\circ 50'$, while for $l > 102^\circ 45'$, the emission is concentrated along $b \simeq -0^\circ 35'$. The whole structure will be referred as cloud A. Two smaller molecular clouds are detected at $(l, b) \sim (103^\circ 2', -0^\circ 30')$ (indicated in the figure as cloud C) and at $(l, b) \sim (103^\circ 4', -0^\circ 42')$ (cloud B). The emission at 8.3 μm correlates very well with the molecular emission. In particular, the brighter emission regions in the mid infrared coincide with clumps in cloud A [for example at $(l, b) \sim (102^\circ 43', -0^\circ 35')$, $(102^\circ 44', -0^\circ 43')$, $(102^\circ 47', -0^\circ 44')$, and $(103^\circ 5', -0^\circ 42')$]. Cloud B coincides with the strong radio source NVSS J222034+561438, which is an YSO candidate. The comparison of the ionized and molecular gas distributions shows that the molecular emission encircles the brightest optical emission regions analyzed by Sánchez-Monge et al. (2008).

The lower panel displays the molecular emission between -53.4 to -46.0 km s^{-1} . The CO emission is clumpy, being concentrated in clouds D, E and F. Cloud C is also detected in this velocity range. Cloud E overlaps a section of


Table 4. IR filaments and point sources.

Source	Fluxes [Jy]								T_d (K)	$L_{IRAS}(L_\odot)$
	8.3 μm	12.1 μm	14.7 μm	21.3 μm	12 μm	25 μm	60 μm	100 μm		
f1	46 $\pm$ 1	—	—	—	72 $\pm$ 4	192 $\pm$ 24	1370 $\pm$ 130	2765 $\pm$ 145	32 $\pm$ 7	43800
f2	16.0 $\pm$ 3.5	—	—	—	35 $\pm$ 6	77 $\pm$ 11	484 $\pm$ 51	1201 $\pm$ 160	29 $\pm$ 6	16900
f3	1.8 $\pm$ 0.3	—	—	—	4.9 $\pm$ 0.5	32 $\pm$ 11	133 $\pm$ 40	343 $\pm$ 37	32 $\pm$ 6	5000
ps1	1.45	1.85	0.97	2.14	3.03	4.28	74.1	274	28 $\pm$ 5	2700
ps2	2.22	3.93	3.78	7.02	7.8	16.1	106	318	30 $\pm$ 5	3900
ps3	1.0	1.6	1.40	4.26	7.02	17.2	243	453	36 $\pm$ 8	6800

**Figure 6.** IR spectral energy distribution for the IR filaments. The lines in the diagram corresponding to $f1$, $f2$, and $f3$ show the emission of a blackbody with a $T = 32$, 29 and 32 K, respectively, and a $\nu^{1.5}$ -emissivity law. The boxes denote flux densities in the four IRAS bands, and the circle, that corresponding to the MSX band A.

cloud A and follows the ring-like feature identified at $8.3 \mu\text{m}$. Cloud F coincides with part of the IR filament at $(l, b) \sim (102^\circ 44', -0^\circ 42')$. The section of Cloud D running from $(l, b) = (102^\circ 56', -00^\circ 51')$ to $(l, b) = (103^\circ 02', -00^\circ 44')$ coincides with f3. Cloud D also displays a clump at $(l, b) \sim (103^\circ 05', -0^\circ 44')$, which coincides with a region of high optical extinction and is probably connected with cloud B.

Table 5 gives the physical parameters of each molecular cloud. The masses of the different clouds were estimated by integrating the CO line intensity as $W(\text{CO}) = \int T_{mb} dv = T_{mb} \Delta v$, where T_{mb} is the average main beam brightness temperature of the molecular cloud over the velocity in-

**Figure 7.** Overlay of the averaged ^{12}CO emission distribution for the molecular clouds between -46.0 to -38.6 km s^{-1} (upper panel) and -53.4 to -46.0 km s^{-1} (lower panel) and the MSX band A emission (grayscale). Contours correspond to 3 , 7 , and 15σ . The grayscale goes from 0.1×10^{-7} to $1 \times 10^{-7} \text{ Watts m}^{-2} \text{ ster}^{-1}$. The crosses indicate the positions of the massive stars.

terval in which the cloud is observed. To estimate the H_2 column density $N(\text{H}_2)$, the relation $X = N(\text{H}_2)/W(\text{CO}) = (2.3 \pm 1.2) \times 10^{20} \text{ mol cm}^{-2} \text{ K}^{-1} \text{ km}^{-1} \text{ s}$ (Grenier & Lebrun 1990) was used. The molecular mass was derived from $M(M_\odot) = 4.2 \times 10^{20} N(\text{H}_2) d^2 \Omega$, where d is the distance in pc and Ω is the solid angle in steradians. Errors in T_{mb} and in $N(\text{H}_2)$ arise in background uncertainties, while errors in masses come from background and distance uncertainties. The adopted distance is $3.5 \pm 1.0 \text{ kpc}$.

To determine the volume density, we assumed a spheri-

Table 5. Physical parameters of the molecular clouds.

	Clouds					
	A	B	C	D	E	F
Average T_{mb} (K)	1.30±0.30	0.90±0.15	0.60±0.15	1.20±0.30	1.60±0.15	0.70±0.15
Velocity range (km s ⁻¹)	-38.6,-46.0	-38.6,-46.0	-38.6,-53.4	-46.0,-53.4	-46.0,-53.4	-46.0,-53.4
$N(H_2)$ column density (10 ²⁰ cm ⁻²)	6.50±1.60	3.30±0.60	4.20±1.05	5.90±2.40	9.0±0.90	1.95±0.20
H_2 mass (10 ³ M _⊙)	4.25±1.90	0.80±0.35	0.04±0.02	1.30±0.60	1.55±0.70	0.95±0.45
Volume density (cm ⁻³)	37±22	2150±1290	110±65	95±55	115±70	320±190
$N(HI)$ column density (10 ²¹ cm ⁻²)	3.7±0.9	3.5±0.5	3.7±0.9	3.3±0.8	3.4±0.3	3.3±0.6
Visual absorption A_v (mag)	2.5±0.8	2.2±0.5	2.8±0.7	2.4±0.7	2.8±0.6	2.7±0.5

cal geometry for all the clouds, except for Cloud A, for which an elongated geometry was adopted.

Figure 8 shows two velocity-position maps of the CO emission. We have plotted Galactic longitude in the y-axis since the molecular morphology is mostly elongated in this direction. The map in the upper panel is the result of integrating within $\Delta b = (-0^\circ 55' 30'', -0^\circ 39' 54'')$, while the map in the bottom panel was integrated within $\Delta b = (-0^\circ 39' 54'', -0^\circ 29' 06'')$. The upper panel shows two clouds at $l \sim 102^\circ 45'$ and $l \sim 103^\circ 00'$, having velocities in the range $\Delta v \sim (-40, -45)$ km s⁻¹ and $(-45, -50)$ km s⁻¹, respectively (two small cloudlets are also detected at $v \simeq -30$ km s⁻¹ at $l \sim 102^\circ 50'$ and $l \sim 103^\circ 03'$). Clouds F and D are clearly identified. The bottom panel shows cloud A extending from $-102^\circ 35'$ to $103^\circ 13'$. The velocity range goes from -40 km s⁻¹ at $l \sim 103^\circ 10'$ to -50 km s⁻¹ at $l \sim 102^\circ 35'$. The image shows some small cloudlets at $l \sim 103^\circ 23'$ at $v \simeq -38$ km s⁻¹, $l < 102^\circ 40'$ with velocities in the range ~ -8 to -5 km s⁻¹, and at $l > 103^\circ 25'$ at $v \simeq 0$ km s⁻¹. Molecular gas with velocities higher than -10 km s⁻¹ is unconnected to Sh2-132.

The agreement in velocity between the ionized and molecular gas and the fact that the molecular emission borders the brightest ionized regions indicate that molecular gas with velocities between -38 and -54 km s⁻¹ is associated with Sh2-132 and the massive stars in the region.

We adopt -45 km s⁻¹ as the systemic velocity of the molecular gas. Taking into account the non-circular motions in this section of the Galaxy shown by Brand & Blitz (1993), a kinematical distance $d_k = 3.0 \pm 1.0$ kpc can be estimated, compatible with the distance adopted in Sect.1.

3.4 The distribution of the HI emission

The upper panel of Fig.9 displays the HI emission distribution in the velocity range where the CO emission is present, i.e. between -31.1 and -58.4 km s⁻¹. The white circle has the dimensions of the HII region. Neither a region lacking HI gas coincident in position with Sh2-132 as a whole nor an HI shell surrounding it are evident in the image, although the small cavity near $(l, b) \simeq (102^\circ 58', -0^\circ 40')$ is probably linked to shell A.

However, a gradient in the HI emission distribution is evident for $b < -0^\circ 10'$. The lower panel displays the emission corresponding to $l = 102^\circ 50'$ from $b = 0^\circ 00'$ to $-1^\circ 30'$ from the image in the upper panel. T_b and $N(HI)$ are shown on left and right y-axes, respectively. The image shows that the HI brightness temperature is $T_b \simeq 85$ K at $b \simeq -0^\circ 15'$,

while 65% this value is detected at $b \simeq -1^\circ 30'$. The **fitted curve**, showed as a dashed line, confirms the presence of a T_b -gradient.

The fitting curve was used to estimate the contribution of the HI gas to the visual absorption A_v . We calculated the visual absorption A_v due to the molecular and neutral hydrogen related to the HII region by using the relation $(N(HI) + 2N(H_2))/E(B-V) = 5.8 \times 10^{21}$ atoms cm⁻² mag⁻¹ (Bohlin et al. 1978), where $N(HI)$ and $N(H_2)$ are the HI and H₂ column densities, respectively, while $E(B-V)$ is the color excess. We adopted $E(B-V) = A_v/3.05$ and read off $N(HI)(b)$ from the diagram. The estimated HI column densities, as well as the derived A_v -values for each molecular cloud are included in Table 5.

4 PHOTODISSOCIATED REGIONS

The presence of PAH emission at the interphase between the ionized and molecular materia, e.i. at the border of many HII regions, suggests that molecular gas is being photodissociated by the UV photons emitted by the massive stars. In the following subsections we analyze the presence of PDRs in two selected areas within the HII region, one around WR 153ab and BD+55°2722, linked to shell B, and the other around LS+55°39, linked to shell A.

4.1 Region near WR 153ab and BD+55°2722

The upper panel of Figure 10 displays an overlay of the DSS-R image and the CO emission distribution for a small region of Sh2-132 near WR 153ab and BD+55°2722. Thin contours correspond to Cloud A with velocities in the range -38.6 to -46.0 km s⁻¹, while thick ones delineate Clouds E and F having velocities between -46.0 and -53.4 km s⁻¹. The crosses mark the position of the stars.

The morphological agreement between the optical emission and the inner borders of the molecular clouds is excellent, particularly for the 4' optical filament at $(l, b) \sim (102^\circ 46' 30'', -0^\circ 43')$, where the contour corresponding to 2.25 K closely follows the optical emission. Note also that diffuse emission is also encircled by molecular emission. The existence of emission at $8.3 \mu\text{m}$ bordering the ionized region (see Fig. 7) strongly indicates the presence of a PDR.

To confirm the existence of a PDR in this region we compared the distribution of the ionized and molecular gas, and that of PAHs. After convolving the DSSR and MSX




Figure 8. *Upper panel:* Averaged ^{12}CO velocity distribution for $\Delta b = (-55'30'', -39'54'')$. The grayscale goes from 0.15 to 3 K. Contours are 0.45, 0.75 and 1.05 K. *Bottom panel:* Averaged ^{12}CO velocity distribution between $\Delta b = (-39'54'', -29'06'')$. Contours and grayscale are the same as in the upper panel.

band A images to the CO beam ($46''$), we averaged the optical, molecular and band A emissions in concentric rings from the position of the WR star within an angle of 300° from $(l, b) \sim (102^\circ 52', -0^\circ 38')$ clockwise up to $(l, b) \sim (102^\circ 49', -0^\circ 33')$. The result of this average is displayed in the bottom panel of the figure, where intensities in the different bands are normalized to the maximum value of each band. The values in the abscissae indicates distance from the WR star.




Figure 9. *Upper panel:* Average HI emission distribution between -31.1 and -58.4 km s^{-1} . The grayscale goes from 50 to 93 K, while contours correspond to 60, 70, 80, and 90 K. The white circle has the dimensions of the HII region Sh2-132. *Bottom panel:* Diagram showing T_b vs. b from $b = -90'$ to $0'$ for $l = 102^\circ 50'$ from the image above. The dashed line shows the best fit to the data.

The diagram reveals a clear stratification, with the optical emission peaking near the stellar position, and the emission at $8.3 \mu\text{m}$ peaking $3'$ far from the star. The molecular emission increases up to $6'$ from the star, where the maximum is present. The observed distribution is typical of molecular clouds exposed to UV radiation, which photodissociates them (van der Werf et al. 1996).

4.2 Region near LS+55°39

The region around LS+55°39 is particularly interesting. The upper panel of Fig. 11 shows that the star is close to a bright optical filament of $\sim 2'$ in size. The fact that the optical filament partially follows the border of cloud A indicates that the UV photons of the star are photodissociating and ionizing the dense gas. The high extinction region at $(l, b) \sim (102^\circ 57', -0^\circ 40')$ coincides with a section of cloud D.

With the aim of investigating the presence of a PDR near the optical filament, we plotted the normalized intensity of optical, MSX band A, and CO emissions as a function of the distance to LS+55°39. The plotted emission corresponds to the region indicated by the black arrow in the upper panel. As for the region around WR 153ab and BD+55°2722, the optical emission peaks closer to the ex-




Figure 10. *Upper panel:* Overlay of the DSS-R image and the CO emission in the region near WR 153 and LS+55°2722, indicated as crosses. Thin lines delineate Cloud A and thick lines, clouds E and F. *Bottom panel:* Normalized intensities of the optical, MSX band A, and CO emissions in concentric rings as a function of the distance to the WR star.

citation source than the emission in MSX band A and the molecular emission, revealing the presence of a PDR.

Taking into account that LS+55°39 is a B0V star with $\log Q(H^0) = 46.23$ (Vacca et al. 1996), and assuming that it is immersed in an ionized interstellar medium with a mean electron density $n_e = 20 \text{ cm}^{-3}$, we can estimate the Strömgen radius r_s of the HII region created by the star from $\frac{3Q(H^0)}{4\pi\alpha_B n_e^2} = r_s^3$ (Mezger & Henderson 1967), where $\alpha_B = 2 \times 10^{-13} \text{ cm}^3 \text{ s}^{-1}$. The result is $r_s = 5.3 \text{ pc}$. Bearing in mind that the angular distance from the star to the optical filament is $2.2'$, or $2.25 \pm 0.7 \text{ pc}$ at $3.5 \pm 1.0 \text{ kpc}$, we can conclude that the B-type star is responsible for ionizing the gas in the region and contributing to the photodissociation the the molecular gas.




Figure 11. *Upper panel:* Overlay of the DSS-R image and the CO emission in the region near LS+55°39, indicated by a cross. Thin lines delineate Clouds A and D. *Bottom panel:* Normalized intensities of the optical, MSX band A, and CO emissions in concentric rings as a function of the distance to the B star.

5 SCENARIO AND CONCLUSIONS

In this paper we have analyzed the distribution of the ionized, neutral atomic and molecular gas and that of the interstellar dust in Sh2-132.

This study revealed the presence of a number of molecular clouds encircling the ionized nebula, with a mass of $9000 M_\odot$. The velocity of the molecular gas, which is in the range from -38.6 to -53.4 km s^{-1} coincides with the velocities of the $H\alpha$ line, and He and H radio recombination lines, strongly indicating that the ionized gas is associated with the nebula and interacting with the ionized gas.

The ring-like appearance of the gas and dust distribution in the environs of the WR star suggests the action of stellar winds, that sweep up and compress the gas. The mechanical energy E_w released by the multiple system WR 153ab into the ISM can be roughly estimated taking into account the two WR components of the binary system during the duration of the WR phase of a massive star, the previous main sequence phase of the current WR stars, and

the two O-star components. We adopt a mass loss rate $\log(\dot{M}) = -6.64$ and a terminal velocity $v_\infty = 1950 \text{ km s}^{-1}$, corresponding to a O8.5V star (Smith et al. 2002) as parameters of the stellar wind of the O-type components. For the WR components we adopt a conservative mass loss rate $\dot{M} = 5 \times 10^{-6} M_\odot \text{ yr}^{-1}$ and a terminal velocity $v_\infty = 2000 \text{ km s}^{-1}$. Adopting lifetimes of $3 \times 10^6 \text{ yr}$ and $0.5 \times 10^6 \text{ yr}$ for the O and WR phases of the stars, respectively (Conti & Vacca 1990, Meynet & Maeder 2005), E_w results to be $\sim 7.2 \times 10^{50} \text{ erg}$.

Considering the molecular clouds E, F and a section of cloud A, and the dust and ionized components, with an expansion velocity of 7.4 km s^{-1} , the kinetic energy of shell B is $\sim 2.80 \times 10^{48} \text{ erg}$. This result indicates that the multiple system can shape shell B.

The emission at $8.3 \mu\text{m}$ appears concentrated in a ring-like structure bordering the brightest sections of the nebula, indicating that molecular material is being photodissociated at the interface between the ionized and molecular gas. This configuration resembles the IR bubbles analyzed by Watson et al. (2008). To confirm the presence of photodissociation regions we have analyzed in some detail two regions within the complex. One in Shell B around WR 153ab and BD+55°2722, and the other in Shell A around LS+55°39. The relative positions of the ionized gas, PAHs, and molecular gas in these regions show the stratification typical in PDRs. Thus, the distribution of H_2 plays a main role in shaping the HII region.

The emission distribution at $60 \mu\text{m}$ shows the same ring-like structure, along with emission inside the ring. The presence of emission in the far IR inside the HII region, which originates from large dust grains, is compatible with the existence of PDRs since PAHs are destroyed inside HII regions, while large grains are not.

The number of UV photons emitted by the massive stars in the region is seven times higher than the UV photons used to ionize the gas, indicating that the massive stars are responsible for the ionization of the region. Clearly, a large fraction of the available UV photons are used in the heating of the interstellar gas and in the photodissociation of the dense material.

Different studies of HII regions such as those towards RCW 79 (Zavagno et al. 2006), RCW 82 (Pomarés et al. 2009), and Sh2-212 (Deharveng et al 2008) show how the interaction between the massive stars with the surrounding molecular material induces star forming regions through the “collect and collapse” process (Elmegreen and Lada 1977) in the borders of these HII regions. For example, the paper on RCW 79 (Zavagno et al. 2006) reveals the formation of YSOs inside pre-existing molecular cores located at the border of the ionized region. By using the analytical model by Whitworth et al. (1994) to determine the time at which the fragmentation occurs and the size of the structure at that time, they conclude that the collect and collapse process may not be the only one at work. Sh2-212 (Deharveng et al. 2008) and Sh2-104 (Deharveng et al. 2003) represent good examples of stellar formation in which the molecular clouds surround the HII regions and the PAH emission is located at the interphase between the ionized and the molecular gases. In particular, Sh2-104 is encircled by a molecular ring showing four dense molecular clouds containing many dense cores. This configuration is a strong evidence in favour of the trig-

gering star formation process described by the “collect and collapse” model. In all of the preceding examples, we have an HII region generated by an O-type star or a cluster of massive stars.

The region around Shell B presents a similar interstellar scenario as the preceding examples, e.i an optical shell bounded by a strong arc-like feature of PAH emission which are surrounding by molecular material (Clouds A, E, and F). In such scenario, star formation can be induced by the action of the massive stars through the ionizing flux and stellar winds. Indeed, the presence of YSOs towards the molecular clouds linked to Sh2-132 indicates that star formation is very active in this region of the Galaxy (Vasquez et al. in preparation).

ACKNOWLEDGEMENTS

We acknowledge the anonymous referee of her/his comments. This project was partially financed by the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET) of Argentina under projects PIP 112-200801-02488 and PIP 112-200801-01299, Universidad Nacional de La Plata (UNLP) under project 11/G093, Universidad de Buenos Aires under project UBACyT X482, and Agencia Nacional de Promoción Científica y Tecnológica (ANPCYT) under projects PICT 00812 and 2007-00902. The Digitized Sky Survey (DSS) was produced at the Space Telescope Science Institute under US Government grant NAGW-2166. This work was partly (S.P.) supported by the Natural Sciences and Engineering Research Council of Canada (NSERC) and the Fonds FQRNT of Québec. The DRAO Synthesis Telescope is operated as a national facility by the National Research Council of Canada. The CGPS is a Canadian project with international partners and is supported by grants from NSERC. Data from the CGPS is publicly available through the facilities of the Canadian Astronomy Data Centre (<http://cad.c.hia.nrc.ca>) operated by the Herzberg Institute of Astrophysics, NRC.

REFERENCES

- Bakes, E.L.O., Tielens, A.G.G.M. 1998, ApJ, 499, 258
- Bohlin, R.C., Savage, B.D., Drake, J.F. 1978, ApJ, 224, 132
- Brand J., Blitz L. 1993, A&A, 275, 67
- Cappa C.E., Vasquez J., Pineault, S., Cichowski S. 20010, MNRAS, in press
- Chaisson, E.J. 1976 in *Frontiers of Astrophysics*, edited by E.H. Avrett, Cambridge: Harvard Univ. Press, 259
- Chu, Y.-H., 1981, ApJ, 249, 195
- Chu, Y.-H., Treffers R.R. 1981, ApJ, 250, 615
- Chu, Y.-H., Treffers R.R., Kwitter, K.B. 1983, ApJS, 53, 937
- Churchwell E., Walmsley C.M., 1973, A&A, 23, 117
- Condon, J.J., Cotton, W.D., Greisen, E.W., Yin, Q.F., Perley, R.A., Taylor, G.B., Broderick, J.J. 1998, AJ, 115, 1693
- Conti P.S., Vacca W.D. 1990, AJ, 100, 431
- Deharveng L., Lefloch B., Zavagno A., Caplan J., Whitworth A. P., Nadeau D., Martn S., 2003, A&A, 408, 25

- Deharveng L., Zavagno A., Caplan J., 2005, *A&A*, 433, 565
- Deharveng L., Lefloch B., Kurtz S., Nadeau D., Pomarés M., Caplan J., Zavagno A., 2008, *A&A*, 482, 585
- Douglas, J.N., Bash F.N., Bozayan F.A., Torrence G.W., Wolfe C. 1996, *AJ*, 111, 1945
- Elmegreen B.G., Lada C.J., 1977, *ApJ*, 214, 725
- Esteban C. Rosado M. 1995, *A&A*, 304, 491
- Fich M., Blitz M., Stark A.A. 1989, *ApJ*, 342, 272
- Foster, T., Routledge, D. 2003, *ApJ*, 598, 1005
- Georgelin Y. M., Georgelin Y. P. 1976, *A&A*, 49, 57
- Gower J.F.R., Scott P.F., Wills D. 1967, *MNRAS*, 71, 49
- Grenier I.A., Lebrun F. 1990, *ApJ*, 360, 129
- Habing H.J. 1968, *Bull. Astron. Inst. Astrophys.*, 204, 253
- Harten R. H., Felli M., Tofani G. 1978, *A&A*, 70, 205
- Heckathorn J. N., Bruhweiler F. C., Gull T. R. 1982, *ApJ*, 252, 230
- Hogerheijde M.R., Jansen D.J., van Dishoeck E.F. 1995, *AA*, 294, 792
- Junkes N., Fuerst E., Reich W. 1992, *A&A*, 261, 289
- Kaufman M.J., Wolfire M.g. Hollembach D.J., Luhman M.L. 1999, *ApJ*, 527, 795
- Landolt-Rörmstein, 1982, *Numerical Data and Functional Relationships in Science and Technology*, Group VI: Astronomy, Astrophysics and Space Research, Volumen 2: Astronomy and Astrophysics, edited by Hellwege K.H., Lumsden S.L., Hoare M.G., Oudmaijer R.D., Richards D. 2002, *MNRAS*, 336, 621
- Massey, P. 1981, *PASP*, 93, 549
- Meynet G., Maeder A. 2003, *A&A*, 404, 975
- Mezger P.G, Henderson A.P. 1967, *ApJ*, 147, 471
- Miller G.,J., Chu Y.-H. 1993, *ApJS*, 85, 137
- Nugis T., Lamers H. J. G. L. M., 2000, *A&A*, 360, 227
- Pomarés M., Zavagno A., Deharveng L., Cunningham M., Jones P., Kurtz S., Russeil D., Caplan J., Comern F., 2009, *A&A*, 494, 987
- Panov K.P., Seggewiss W., 1990, *A&A*, 227, 117
- Povich M.S., Stone J.M., Churchwell E., Zweibel E.G., Wolfire M.G., Babler B.L., Indebetouw R., Meade M.R., Whitney B.A., 2007, *ApJ*, 660, 346
- Price S.D. Egan M.P., Carey S.J., Mizuno D.R., Kuchar T.A., 2001, *AJ*, 121, 2819
- Prinja R. K., Barlow M.J., Howarth I.D., 1990, *ApJ*, 361, 607
- Quireza C., Rood R.T., Balser D.S., Bania T. M. 2006, *ApJS*, 165, 338
- Quireza C., Rood R.T., Bania T. M., Balser, D.S., Maciel, W.J. 2006, *ApJ*, 653, 1226
- Reich W., Fuerst E., Reich P., Reif K., 1990, *A&A*, 85, 633
- Reynolds R. J., 1988, *ApJ*, 333, 341
- Sánchez-Monge A., Palau A., Estalella R., Beltrán M.T., Girart J.M., 2008, *A&A*, 485, 497
- Sharpless S., 1959, *ApJS*, 4, 257
- Scoville N.Z., Kwan J. 1976, *ApJ*, 206, 718
- Smith L.J., Norris R.P.F., Crowther P.A., 2002, *MNRAS*, 337, 1309
- Smith L.F., Shara M.M., Moffat A.F.J., 1998, *MNRAS*, 281, 163
- Sota A., Maz-Apellniz J., Walborn N.R., Shida R.Y. 2008, *RMxAC*, 33, 56
- Sternberg, A. and Dalgarno, A., *ApJS*, 1995, 99, 565
- Sternberg, A. and Dalgarno, A., *ApJ*, 1989, 338, 197
- Schwartz P.R., *ApJ*, 1982, 252, 589
- Taylor, A.R., Gibson S.J., Peracaula M., Martin P.G., Landecker T.L., Brunt C.M., Dewdney P.E., Dougherty S.M., Gray A.D., Higgs L.A., Kerton C.R., Knee L.B.G., Kothes R., Purton C.R., Uyaniker B., Wallace B.J., Willis A.G., & Durand D., 2003, *AJ*, 125, 3145
- Tielens A.G.G.M., Hollenbach, D. 1985, *ApJ*, 291, 722
- Vacca W.D., Garmany C.D., Shull J.M., 1996, *ApJ*, 460, 914
- van der Hucht K.A., 2001, *New Astronomy. Rev.*, 45, 135
- Waldram E.M., Yates J.A., Riley J.M., Warner P.J., 1996, *MNRAS*, 282, 779
- Wegner W., 1994, *MNRAS*, 270, 229
- Watson C., Povich M.S., Churchwell, E.B., Babler B.L., Chunev G. et al. 2008, *ApJ*, 681, 1341
- Whittet D.C.B., from the book “Dust in the galactic environment”, Cambridge University Press
- Whitworth A.P., Bhattal A.S., Chapman S.J., Disney M.J., Turner J.A., 1994, *A&A*, 290, 421
- van der Werf P.P., Stutzki J., Sternberg A., Krabbe A. 1996, *AA*, 313, 633
- Zavagno A., Deharveng L., Comern F., Brand J., Massi F., Caplan J., Russeil D., 2006, *A&A*, 446, 171